



Available at
WWW.MATHEMATICSWEB.ORG
 POWERED BY SCIENCE @ DIRECT®

**JOURNAL OF
 Algebra**

Journal of Algebra 264 (2003) 251–261

www.elsevier.com/locate/jalgebra

CS matrix rings over local rings

K.I. Beidar,^a S.K. Jain,^b Pramod Kanwar,^c and J.B. Srivastava^d

^a Department of Mathematics, National Cheng-Kung University, Tainan, Taiwan

^b Department of Mathematics, Ohio University, Athens, OH 45701, USA

^c Department of Mathematics, Ohio University–Zanesville, Zanesville, OH 43701, USA

^d Department of Mathematics, Indian Institute of Technology, Delhi 110016, India

Received 5 May 2002

Communicated by Kent R. Fuller

Abstract

Complete characterization of CS matrix rings $M_n(R)$, $n > 1$, over local rings R is obtained. Application to group algebras is derived as a particular case of the main result.

© 2003 Elsevier Science (USA). All rights reserved.

Keywords: CS-rings; Finitely \sum -CS modules; Local rings; Semiperfect rings; Locally finite groups; Nilpotent groups

1. Introduction

A ring R is called a right CS-ring if every essentially closed right ideal is a direct summand of R . Such rings have been studied by several authors (cf. [2–9]). It is known that if R is a commutative integral domain then $M_2(R)$ is a right CS-ring if and only if R is a Prüfer domain [5, Corollary 12.10] and that if R is a local (noncommutative) domain then $M_n(R)$, $n > 1$, is a right CS-ring if and only if R is a valuation domain [1, Lemma 3.6]. In this paper we first show that the $n \times n$ matrix ring ($n > 1$) over a local ring R is right CS if and only if R is right uniform and for every right ideal K of R and for every R -homomorphism $f: K \rightarrow R$ there exists $u \in R$ such that either $f = l_u$ or $l_u f = I_K$, where l_u is the left multiplication by u and I_K is the identity map on K (Theorem 3.5). If, in addition, the radical of R coincides with the right singular ideal, then $M_n(R)$, $n > 1$, is a right CS-ring if and only if R is a right selfinjective ring (Theorem 3.6).

E-mail addresses: kbeidar@yahoo.com (K.I. Beidar), jain@math.ohiou.edu (S.K. Jain), pkanwar@math.ohiou.edu (P. Kanwar), jbsrivas@maths.iitd.ernet.in (J.B. Srivastava).

Theorem 3.9 shows that if R is a commutative noetherian local ring then $M_n(R)$, $n > 1$, is a right CS-ring if and only if the classical quotient ring, $Q_{cl}(R)$, is a local QF-ring such that for all $a \in Q_{cl}(R)$ either $a \in R$ or a is invertible and $a^{-1} \in R$. Lemma 3.3, which is also of independent interest, describes all uniform summands of the right R -module R^n and plays an important role in the proof of our main result. In Section 4 we apply our machinery, developed in Section 3, to the local CS group algebras and also to semiperfect group algebras of nilpotent groups. For local group algebra KG of any group G it is shown that $M_n(KG)$, $n > 1$, is a right CS-ring if and only if $\text{char}(K) = p$ and G is a finite p -group.

2. Notation and definitions

Throughout this paper, unless otherwise stated, all rings have unity and all modules are right unital. For any two right R -modules M and N , M is said to be N -injective if for any submodule L of N and any R -homomorphism $\phi: L \rightarrow M$ there exists an R -homomorphism $\psi: N \rightarrow M$ such that $\psi|_L = \phi$. A right R -module M is said to be injective if M is N -injective for all right R -modules N . A submodule K of a right R -module M is said to be essential in M , denoted by $K \subseteq_e M$, if for any nonzero submodule L of M , $K \cap L \neq 0$. M is called a CS (or extending) module if every submodule of M is essential in a direct summand of M , equivalently, if every closed submodule of M is a direct summand of M . M is called finitely \sum -CS if direct sum of finite number of copies of M is CS. M is called CS with respect to uniform submodules if every uniform submodule of M is essential in a direct summand of M , equivalently, if every uniform closed submodule of M is a direct summand of M . M is said to satisfy condition C_3 if for any two summands M_1 and M_2 of M with $M_1 \cap M_2 = 0$, $M_1 \oplus M_2$ is also a summand of M . A CS module is called quasi-continuous if it satisfies C_3 . It is known that if $M \times N$ is quasi-continuous then M and N are injective relative to each other.

A ring R is said to be right CS (or CS with respect to uniform right ideals) if the right R -module R is CS (resp. CS with respect to its uniform right R -submodules). R is called right selfinjective if R_R is injective. R is called a right valuation ring if for any two right ideals I and J either $I \subset J$ or $J \subset I$. Let S be an overring of R . The subset $\{1 = a_1, a_2, \dots, a_n\}$ of S is said to be a normalizing basis of $S_R(RS)$ if $a_i R = R a_i$, $1 \leq i \leq n$. S is called R -projective if for any S -module M and for any S -submodule N of M , if N is an R -summand of M then it is also an S -summand of M . For a ring R , $J(R)$ will denote the Jacobson radical of R and $Z_r(R)$, the right singular ideal $\{r \in R \mid rI = 0 \text{ for some essential right ideal } I \text{ of } R\}$ of R . For a nonempty subset X of a ring R , $\text{r.ann}_R(X)$ ($\text{l.ann}_R(X)$) will denote the right (left) annihilator of X in R . If X is the singleton $\{a\}$ then we write $\text{r.ann}_R(X) = \text{r.ann}_R(a)$ ($\text{l.ann}_R(X) = \text{l.ann}_R(a)$). For an element a of R , l_a will denote left multiplication by a .

A group G is called locally finite if every finitely generated subgroup of G is finite. For a group G , $O_p(G)$ will denote the maximal normal p -subgroup and $\omega(RG)$ will denote the augmentation ideal of the group ring RG . If H is a subgroup of G , we will write $\omega(H)$ to denote $\omega(RH)RG$.

3. Main results

Lemma 3.1 [5, Corollary 7.8]. *A right module over a ring R with finite uniform dimension is CS if and only if it is CS with respect to uniform submodules.*

Lemma 3.2 [5, Lemma 12.8]. *The matrix ring $M_n(R)$ over a ring R is right CS if and only if R^n is a CS-module as a right R -module.*

Lemma 3.3. *Suppose R is a local right CS-ring. A uniform right R -submodule U of R^n is a summand of R^n if and only if $U = (a_1, a_2, \dots, a_n)R$, where some $a_i = 1$.*

Proof. Since R is a local right CS-ring, R is a uniform right R -module. Let U be a uniform summand of the right R -module R^n . Then $R^n = U \oplus K$ for some right R -submodule K of R^n . Since R is local right uniform and U is uniform, by Krull–Schmidt Theorem, $U \simeq R$ as right R -modules. Let α be the isomorphism from R to U such that $1 \rightarrow (a_1, a_2, \dots, a_n) \in R^n$. Then $U = (a_1, a_2, \dots, a_n)R$. Consider the R -isomorphism $f: U \rightarrow R$ where $f = \alpha^{-1}$. Extend f to f^* from $U \oplus K = R^n$ to R by setting $f^* = f$ on U , and $f^* = 0$ on K . Now every homomorphism from R^n to R can be represented by a $n \times 1$ matrix with entries in R . Let

$$f^* = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}.$$

Since $(a_1, a_2, \dots, a_n) \in U$ is the preimage of $1 \in R$ under f ,

$$(a_1, a_2, \dots, a_n) \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = 1,$$

that is, $a_1x_1 + a_2x_2 + \dots + a_nx_n = 1$. If a_1, a_2, \dots, a_n are all in $J(R)$ then $1 \in J(R)$, a contradiction. Hence there exists i , $1 \leq i \leq n$, such that a_i is a unit. But then $U = (b_1, b_2, \dots, b_{i-1}, 1, b_{i+1}, \dots, b_n)R$ where each $b_j = a_j a_i^{-1}$ as desired. Conversely, any right R -submodule U of the form $(a_1, a_2, \dots, a_{i-1}, 1, a_{i+1}, \dots, a_n)R$ is a summand of R^n because

$$(a_1, a_2, \dots, a_{i-1}, 1, a_{i+1}, \dots, a_n)R \oplus \left(\bigoplus_{\substack{j=1 \\ j \neq i}}^n e_j R \right) = R^n,$$

where $e_j \in R^n$ is the row vector all of whose entries are 0 except the j th entry which is 1. \square

Next we give a simple fact regarding finite linearly preordered sets.

Lemma 3.4. *Let (S, \geq) be a finite set with linear preorder \geq . Then there exists $x \in S$ such that $x \geq s$ for all $s \in S$.*

Proof. The proof follows by induction. \square

Theorem 3.5. *The following are equivalent for a local ring R .*

- (1) $M_n(R)$, $n > 1$, is a right CS-ring.
- (2) $M_2(R)$ is a right CS-ring.
- (3) R is right uniform, and for every right ideal K of R and for every R -homomorphism $f: K \rightarrow R$ there exists $u \in R$ such that either $f = l_u$ or $l_u f = I_K$.

Proof. (1) \Rightarrow (2) is obvious.

(2) \Rightarrow (3) Since $M_2(R)$ is right CS, $R \times R$ is a CS right R -module. Thus R is right CS and hence right uniform. Let K be right ideal of R and let $f: K \rightarrow R$. Let $U = \{(x, f(x)) \mid x \in K\}$. Then $U_R \simeq K_R$. Thus U is uniform. Since $R \times R$ is a CS right R -module, U is essential in a summand S of $R \times R$. By Lemma 3.3, $S = (1, b)R$ or $(a, 1)R$. If $U \subseteq_e (1, b)R$ then for every $x \in K$, there exist $r \in R$ such that $(x, f(x)) = (1, b)r$. Thus $x = r$ and $f(x) = br$. It follows that $f = l_b$. If $S \subseteq_e (a, 1)R$ then for every $x \in K$ there exist $r \in R$ such that $(x, f(x)) = (a, 1)r$. Thus $x = ar$ and $f(x) = r$. Hence $x = af(x) = l_a f(x)$ for every $x \in K$, that is, $l_a f = I_K$. This proves (3).

(3) \Rightarrow (1) By Lemma 3.2 it is sufficient to prove that R^n is CS as a right R -module. Since R is right uniform, by Lemma 3.1, we only need to consider uniform right R -submodules of R^n . Let U be a uniform right R -submodule of R^n . Let π_i , be the canonical projection of R^n onto i th direct summand, let $K_i = \pi_i(U)$ and let f_i be the restriction of π_i onto U , $i = 1, 2, \dots, n$. Clearly,

$$U = \{(f_1(x), f_2(x), \dots, f_n(x)) \mid x \in U\} \quad (1)$$

and $\bigcap_{i=1}^n \ker(f_i) = 0$. Since U is uniform, there exists $1 \leq i \leq n$ such that $\ker(f_i) = 0$. Obviously f_i is an isomorphism. We may assume without loss of generality that there exists a positive integer $k \leq n$ such that f_1, f_2, \dots, f_k are isomorphisms whereas $\ker(f_j) \neq 0$ for all $j = k+1, k+2, \dots, n$. Given $1 \leq i, j \leq k$, $f_i f_j^{-1}: K_j \rightarrow K_i$ is an isomorphism of R -modules. By our assumption there exists $a \in R$ such that either $f_i f_j^{-1} = l_a$ or $l_a f_i f_j^{-1} = I_{K_j}$. Therefore either $f_i = l_a f_j$ or $f_j = l_a f_i$. Now we introduce a linear preorder on $\{1, 2, \dots, k\}$ as follows. We set $i \geq j$ if there exists $a \in R$ such that $f_j = l_a f_i$. Obviously this binary relation \geq is transitive and reflexive since $f_i = l_1 f_i$. Therefore it is a linear preorder relation. By Lemma 3.4, the set $\{1, 2, \dots, k\}$ has a maximum element. Let i be the maximum element of $\{1, 2, \dots, k\}$. We may assume without loss of generality that $i = 1$. Therefore there exist $a_2, a_3, \dots, a_k \in R$ such that $f_j = l_{a_j} f_1$, $j = 2, 3, \dots, k$. Let $k+1 \leq r \leq n$. By our assumption there exist $a_r \in R$ such that either $f_r f_1^{-1} = l_{a_r}$ or

$l_{a_r} f_r f_1^{-1} = I_{K_1}$. The latter possibility is ruled out by the fact that $\ker(f_r) \neq 0$. We see that there exist $a_2, a_3, \dots, a_n \in R$ such that

$$f_j = l_{a_j} f_1, \quad j = 2, 3, \dots, n. \quad (2)$$

Thus from (1) and (2) we obtain

$$\begin{aligned} U &= \{(f_1(x), f_2(x), \dots, f_n(x)) \mid x \in U\} \\ &= (1, a_2, a_3, \dots, a_n) f_1(U) = (1, a_2, a_3, \dots, a_n) K_1. \end{aligned}$$

Therefore $U \subseteq_e (1, a_2, a_3, \dots, a_n)R$ where $(1, a_2, a_3, \dots, a_n)R$ is a direct summand of R^n by Lemma 3.3 and we are done. \square

Theorem 3.6. *Suppose R is a local ring with $J(R) = Z_r(R)$. Then $M_n(R)$ is right CS for some $n > 1$ if and only if R is right selfinjective.*

Proof. Let $M_n(R)$ be right CS for some $n > 1$. By Lemma 3.2, R^n is CS as a right R -module. Consequently R is right CS and hence right uniform. To prove R is right selfinjective, let K be a nonzero right ideal of R and let $f: K \rightarrow R$ be an R -homomorphism. By Theorem 3.5, there exists $u \in R$ such that either $f = l_u$ or $l_u f = I_K$. If $l_u f = I_K$ then f is a monomorphism and $uf(a) = a$ for every $a \in K$. We show that u is invertible in R for otherwise $u \in J(R) = Z_r(R)$. Thus $\text{r.ann}_R(u)$ is essential in R . Since K is nonzero, $f(K)$ is nonzero. Consequently $\text{r.ann}_R(u) \cap f(K) \neq 0$. Let $0 \neq f(a) \in \text{r.ann}_R(u) \cap f(K)$. Then $a = uf(a) = 0$, a contradiction because $f(a) \neq 0$. Hence $f(a) = u^{-1}a$ for every $a \in K$. Thus $f = l_{u^{-1}}$. This proves the result. \square

Since for a right uniform local ring R with nil radical $J(R) = Z_r(R)$, we have the following corollary.

Corollary 3.7. *Suppose R is a local ring with nil radical. Then $M_n(R)$ is right CS for some $n > 1$ if and only if R is right selfinjective.*

We call a ring R right almost selfinjective if for any right ideal K of R and any R -homomorphism $f: K \rightarrow R$ there exists $a \in R$ such that either $f = l_a$ or $l_a f = I_K$.

We do not know whether for a local ring R , $M_n(R)$ ($n > 1$) being right CS implies that $M_n(R)$ is also left CS. In particular, whether the condition that $M_n(R)$ is a right CS-ring implies that R is right-left uniform and right-left almost selfinjective. Theorem 3.9 characterizes local uniform right-left almost selfinjective rings with acc on right-left annihilators. Before proving the theorem we prove the following lemma.

Lemma 3.8. *Let R be a local right selfinjective ring and let A be a subring of R such that for any $a \in R$ either $a \in A$ or a is invertible with $a^{-1} \in A$. Then we have the following.*

- (1) *A satisfies right-left Ore conditions and R is both right as well as left classical ring of quotients of A .*

- (2) R_A is an injective A -module.
 (3) A is a local right uniform almost right selfinjective ring.

Proof. (1) Let S be the set of all elements of A which are not left or right zero divisors in A and let $a \in S$. We first show that a is neither a left nor a right zero divisor in R . If $ax = 0$ for some $0 \neq x \in R$, then $x \notin A$ and so x is invertible in R forcing $a = 0$, a contradiction. Thus a is not a left zero divisor in R . Similarly a is not a right zero divisor in R . We now claim that a is invertible in R . Consider the right R -homomorphism $f : aR \rightarrow R$ given by the rule $f(ay) = y$ for all $y \in R$. Since R is right selfinjective, there exists $b \in R$ such that $f = l_b$ and so $ba = 1$. Therefore $(1 - ab)a = 0$. Since a is not a right zero divisor, we get $ab = 1$. Thus every element of S is invertible in R . By hypothesis, for every $r \in R$, either $r \in A$, or r is invertible in R and $r^{-1} \in A$. Therefore R is both left and right ring of fractions of A (see [14, p. 50]). It now follows that A satisfies both left and right Ore conditions and R is the two-sided classical ring of quotients of A (see [14, Proposition 1.4, p. 51]).

(2) By [14, Proposition 3.5, p. 57], both ${}_A R$ and R_A are flat modules. Since R_R is injective, R_A is an injective module (see [10, Corollary 3.6A]).

(3) Since R is a local right selfinjective ring, R is right uniform. As R is the classical ring of quotients of A , we conclude that A is also right uniform. To show that A is local, let J be the set of all elements of A which are not invertible in A . By [11, Theorem 19.1], it is enough to show that J is closed under addition. Let $a, b \in J$. Assume that $a + b$ is invertible in A . Set $u = a(a + b)^{-1}$ and $v = b(a + b)^{-1}$. Then $u, v \in A$ and $u + v = 1$. As R is local, we may assume that u is invertible in R . Setting $w = u^{-1}v$, we see that either $w \in A$ or $w^{-1} \in A$. In the former case we have $u^{-1} = 1 - u^{-1}v = 1 - w \in A$ and so $a^{-1} = (a + b)^{-1}u^{-1} \in A$, a contradiction. In the latter case $v^{-1} = 1 - w^{-1} \in A$ forcing $b^{-1} = (a + b)^{-1}v^{-1} \in A$, a contradiction again. Thus A is local ring.

To prove that A is right almost selfinjective, let K be a right ideal of A and let $f : K \rightarrow A$ be a right A -homomorphism. Since R is injective as a right A -module, we may assume that $f : R \rightarrow R$. As R is the classical ring of quotients of A , it is easy to see that f is an endomorphism of right R -modules and so there exists $a \in R$ such that $f = l_a$. If $a \in A$, then there is nothing to prove. If $a \notin A$, then a is invertible and $b = a^{-1} \in A$. Therefore $l_b f = I_K$. This completes the proof. \square

Theorem 3.9. *Let R be a ring. Then the following conditions are equivalent.*

- (1) R is local left (or right) uniform, almost left and right selfinjective, and satisfies acc condition on left and right annihilators.
 (2) R satisfies both left and right Ore conditions. Its two-sided classical ring of quotients $Q = Q_{cl}(R)$ is a local QF-ring and for any $a \in Q$ either $a \in R$ or a is invertible in Q with $a^{-1} \in R$.

Proof. (1) \Rightarrow (2) Let S be the set of all elements of R which are not left zero divisors. We claim that every element of S is not a right zero divisor. Indeed, let $a \in S$. Assume that $x_1 a = 0$ for some $0 \neq x_1 \in R$. Since $r.\text{ann}_R(a) = 0$, the map $f_1 : aR \rightarrow x_1 R$, given by the rule $f_1(ay) = x_1 y$, $y \in R$, is a well-defined homomorphism of right R -modules.

By our assumption there exists $x_2 \in R$ such that either $f_1 = l_{x_2}$ or $l_{x_2}f_1 = I_{aR}$. The latter possibility is ruled out by the fact that $f_1(a^2) = x_1a = 0$. Therefore $f_1 = l_{x_2}$. Hence $x_2a = l_{x_2}(a) = f(a) = x_1 \neq 0$. Also $x_2a^2 = x_1a = 0$. Thus $x_2 \in \text{l.ann}_R(a^2) \setminus \text{l.ann}_R(a)$. Continuing in this fashion, we shall construct a strictly increasing chain

$$\text{l.ann}_R(a) \subset \text{l.ann}_R(a^2) \subset \cdots \subset \text{l.ann}_R(a^n) \subset \cdots,$$

contrary to our assumption. Therefore our claim is established.

Given $a \in S$ and $x \in R$, we claim that there exists $b \in S$ and $y \in R$ such that $bx = ya$. Consider the map $f: aR \rightarrow xR$ given by the rule $f(ay) = xy$, $y \in R$. Obviously f is a well-defined homomorphism of right R -modules. By our assumption there exists $z \in R$ such that either $f = l_z$ or $l_zf = I_{aR}$, that is, either $x = za$ or $a = zx$. In either case our assertion is trivially true.

We note that S satisfies the left Ore condition. By left-right symmetry, S satisfies the right Ore condition as well. Therefore, R has the classical ring of quotients Q .

Again let $a \in S$ and $x \in R$. Then as in the previous paragraph, either $x = za$ or $a = zx$ for some $z \in R$, that is, either $x \in Ra$ or $a \in Rx$. Equivalently, either $xa^{-1} \in R$ or $(xa^{-1})^{-1} = ax^{-1} \in R$. We note that for any $y \in Q$ either $y \in R$ or y is invertible and $y^{-1} \in R$. It now follows that for any set $P \subset Q$, $\text{l.ann}_Q(P) = \text{l.ann}_R(P)$ and $\text{r.ann}_Q(P) = \text{r.ann}_R(P)$. In particular, Q satisfies acc condition on left and right annihilators. Since R is a right uniform ring, Q is also a right uniform ring. Further, since R is a local ring, Q is also local. Moreover, every element of $J(Q)$ is a zero divisor and $J(Q) \subset R$.

We now claim that Q is right selfinjective. To prove the claim, let U be a right ideal of Q and let $f: U \rightarrow Q$ be a right Q -homomorphism. We show that f is given by the left multiplication. If $U = Q$ we are done. So let $U \neq Q$ and so $U \subset J(Q) \subset R$. Since every element of U is right zero divisor, $f(U)$ consists of right zero divisors and so $f(U) \subset J(Q) \subset R$. By our assumption there exists $a \in R$ such that either $f = l_a$ or $l_af = I_U$. In the latter case the uniformity of the ring R implies that a is not a left zero divisor in R and so $a \in S$ is invertible in Q forcing $f = l_{a^{-1}}$. Thus either $f = l_a$ or $f = l_{a^{-1}}$, that is, f is given by the left multiplication, as desired. It now follows that Q is a QF-ring (see [14, Theorem 3.5, p. 277]).

(2) \Rightarrow (1) Since any subring of a QF-ring satisfies the acc condition on left and right annihilators, the result follows from Lemma 3.8. \square

Corollary 3.10. *Let R be a commutative noetherian local ring. Then $M_n(R)$, $n > 1$, is a right CS-ring if and only if the classical quotient ring, $Q_{\text{cl}}(R)$, is local QF such that for all $a \in Q_{\text{cl}}(R)$ either $a \in R$ or a is invertible and $a^{-1} \in R$.*

If R is a local ring then $Z_r(R) \subset J(R)$. If $Z_r(R) = 0$ then R is a domain and it is known that for a local domain R , $M_n(R)$, $n > 1$, is right CS if and only if R is a right and left valuation domain [1, Lemma 3.6]. In case $Z_r(R) = J(R)$ then $M_n(R)$ is right CS for some $n > 1$ if and only if R is right selfinjective (Theorem 3.6). We now provide an example of a local ring R such that $M_n(R)$ is right CS, but R is neither a domain nor right selfinjective.

Example 3.1. Let R be a right and left valuation domain, not necessarily commutative and let D be its right classical ring of quotients. Let

$$T = \left\{ \begin{pmatrix} a & d \\ 0 & a \end{pmatrix} \mid a \in R, d \in D \right\} \subseteq S = \left\{ \begin{pmatrix} a & d \\ 0 & a \end{pmatrix} \mid a, d \in D \right\}.$$

Obviously S is a local QF-ring and T is a subring of S such that for every $x \in S$ either $x \in T$ or $x^{-1} \in T$. According to Lemma 3.8, T is a local right uniform ring such that for any right ideal K of T and any right T -homomorphism $f: K \rightarrow T$ there exists $a \in T$ such that either $f = l_a$ or $l_a f = I_K$. Obviously T is not a domain and T is not right selfinjective because $J(T) \neq Z_r(T)$.

We now give another interesting application of Lemma 3.3.

Theorem 3.11. Suppose R is a local right CS-ring with radical equal to the set of all zero divisors. Then $R \times R$, as a right R -module, satisfies C_3 . In particular, if R is a local right CS-ring with nil radical then $R \times R$, as a right R -module, has C_3 .

Proof. It is sufficient to consider uniform summands of $R \times R$ as a right R -module. By Lemma 3.3, a uniform summand of $R \times R$ is either of the type $(1, a)R$ or $(a, 1)R$. To prove our assertion let S_1 and S_2 be uniform summands of R such that $S_1 \cap S_2 = 0$. Thus there exists a and b in R such that $S_1 = (1, a)R$ or $(a, 1)R$ and $S_2 = (1, b)R$ or $(b, 1)R$. First assume that if S_1 and S_2 have 1's in the same position, say, $S_1 = (1, a)R$ and $S_2 = (1, b)R$. Now if both a and b are in $J(R)$ then $\text{r.ann}_R(a - b) \neq 0$. Hence there exists $r \in R$ such that $(a - b)r = 0$. It follows that $0 \neq (1, a)r = (1, b)r \in S_1 \cap S_2$, a contradiction. Hence one of a and b , say a , is not in $J(R)$. But then $S_1 = (1, a)R = (a^{-1}, 1)R$. We, therefore, only need to consider the case when $S_1 = (a, 1)R$ and $S_2 = (1, b)R$. We now show that in this case $S_1 \cap S_2 = 0$ if and only if $1 - ab$ is invertible. First let $1 - ab$ be invertible and let $(a, 1)r = (1, b)s \in S_1 \cap S_2$. Then $ar = s$ and $r = bs$. Thus $abs = s$, that is, $(1 - ab)s = 0$. Since $1 - ab$ is invertible we get $s = 0$. Thus $S_1 \cap S_2 = 0$. Conversely, if $1 - ab$ is not invertible then $1 - ab \in J(R)$. Thus there exists $0 \neq x \in R$ such that $(1 - ab)x = 0$. Hence $0 \neq (a, 1)bx = (1, b)x \in S_1 \cap S_2$. This proves the claim. Next we show that $S_1 \oplus S_2 = R \times R$. It is sufficient to show that $(1, 0)$ and $(0, 1)$ belong to $S_1 \oplus S_2$. Consider the relation $(1, 0) = (a, 1)x + (1, b)y$. Then $ax + y = 1$ and $x + by = 0$. These equations give $(1 - ab)y = 1$ and $x = -by$. Since $1 - ab$ is invertible, $y = (1 - ab)^{-1}$ and $x = -b(1 - ab)^{-1}$. It follows that $(1, 0) \in S_1 \oplus S_2$. Similarly $(0, 1) \in S_1 \oplus S_2$. This completes the proof. \square

Remark 3.1. We note that for the ring R in Theorem 3.11 if $R \times R$ is also CS as a right R -module then $R \times R$ is quasi-continuous and so R is injective as a right R -module [15, Properties 41.20, p. 367]. This gives an alternative proof of Corollary 3.7.

Remark 3.2. Using an argument similar to the one in Theorem 3.11, it can be proved that if R is a local right CS-ring with nil radical then R^n , as a right R -module, has C_3 on uniform summands.

4. Applications to group algebras

In this section we give applications of the results obtained in the previous section to group algebras.

Lemma 4.1. *Let K be a field and G be any group. If the group algebra KG is local right CS then $\text{char}(K) = p$, G is a locally finite p -group, and the radical of KG is nil.*

Proof. Let KG be local right CS. Then $J(KG) = \varpi(KG)$. By [13, Lemma 1.13, p. 415], $\text{char}(K) = p$ and G is a p -group. Also as KG has no nontrivial idempotents, KG is uniform.

We will prove that G is locally finite. Let $H = \langle h_1, h_2, \dots, h_n \rangle$ be a finitely generated subgroup of G . Since G is a p -group, for each i with $1 \leq i \leq n$, $o(h_i) = p^{k_i}$ for some k_i . For each i , let $u_i = 1 + h_i + h_i^2 + \dots + h_i^{p^{k_i}-1}$. Since $u_i KG \neq 0$ for each i and KG is uniform, $\bigcap_{i=1}^n u_i KG \neq 0$. Let α be a nonzero element of $\bigcap_{i=1}^n u_i KG$. Then $(h_i - 1)\alpha = 0$ for each i . Consequently

$$\left(\sum_{i=1}^n KG(h_i - 1) \right) \alpha = 0.$$

Thus $0 \neq \alpha \in \text{r.ann}(\omega(H))$. Hence H is a finite group, as desired. \square

Theorem 4.2. *Let K be a field and G be any group such that the group algebra KG is local. The matrix ring $M_n(KG)$, $n > 1$, is a right CS-ring if and only if $\text{char}(K) = p$ and G is a finite p -group.*

Proof. The proof follows from Corollary 3.7 once we observe that the radical of KG is nil and that KG is right selfinjective if and only if G is finite. \square

We now consider semiperfect group algebras of nilpotent groups.

Theorem 4.3. *Let K be a field and G be a nilpotent group such that the group algebra KG is semiperfect. Then the following are equivalent.*

- (1) $M_n(KG)$, $n > 1$, is a right CS-ring.
- (2) $M_2(KG)$ is a right CS-ring.
- (3) G is finite.

Proof. We only need to prove (2) \Rightarrow (3). Since G is nilpotent, $J(KG)$ is nilpotent. By [13, Theorem 1.5, p. 409] either $\text{char}(K) = 0$ and G is finite or $\text{char}(K) = p$, G is locally finite, and $[G : O_p(G)] < \infty$. We can assume that p does not divide $[G : O_p(G)]$. For if p divides $[G : O_p(G)]$ then taking the unique Sylow p -subgroup $\frac{N}{O_p(G)}$ of $\frac{G}{O_p(G)}$ we get a normal subgroup N of G such that $[G : N] < \infty$ and we can replace $O_p(G)$ with N . Since $O_p(G)$ is normal in G , KG is $KO_p(G)$ -free with normalizing basis, say $\{1 = a_1, a_2, \dots, a_n\}$.

Also because p does not divide $[G : O_p(G)]$, KG is $KO_p(G)$ -projective [13, Lemma 2.2, p. 274]. Let $S = KG$ and $R = KO_p(G)$. Since $M_2(S) = M_2(KG)$ is right CS, $S^2 = S \times S$ is CS as a right S -module.

We show that R^2 is a CS as a right R -module. Observe that $S^2 = R^2a_1 + R^2a_2 + \cdots + R^2a_n$ and there exist automorphisms σ_i ($1 \leq i \leq n$) of the ring R such that $a_i r = \sigma_i(r) a_i$. Let A be a closed R -submodule of R^2 . First we prove that AS is closed R -submodule of S^2 . Note that $AS = Aa_1 + Aa_2 + \cdots + Aa_n$. Let $x = x_1a_{k_1} + x_2a_{k_2} + \cdots + x_ua_{k_u}$ be in the closure of AS in S^2 where $x \notin AS$. We may assume without loss of generality that each $x_i \notin A$. Now there exists an essential right ideal E of R such that $0 \neq xE \subset AS$. But $xy = x_1\sigma_{k_1}(y)a_{k_1} + x_2\sigma_{k_2}(y)a_{k_2} + \cdots + x_u\sigma_{k_u}(y)a_{k_u}$ for every $y \in E$. Since $0 \neq xE \subset AS$ there exists i such that $0 \neq x_i\sigma_{k_i}(E) \subset A$. Because $\sigma_{k_i}(E)$ is essential right ideal of R and A is closed, $x_i \in A$, a contradiction. Hence AS is closed R -submodule of S^2 . Since S is R -projective, AS is a closed S -submodule of S^2 [12, Proposition 1.1]. Consequently AS is a summand of S^2 . Let $S^2 = AS \oplus B$. Writing A_0 for $Aa_2 + \cdots + Aa_n$ and S_0 for $R^2a_2 + \cdots + R^2a_n$, we have $R^2 \oplus S_0 = A \oplus (A_0 \oplus B)$. It follows that $R^2 = A \oplus ((A_0 \oplus B) \cap R^2)$ proving that A is a summand of R^2 . This proves that R^2 is CS as a right R -module. But then $M_2(R)$ is right CS. Since $R = KO_p(G)$ is local, by Theorem 4.2, $O_p(G)$ is finite. Consequently G is a finite group. \square

Note added in proof

(1) It has been pointed out to us that Theorem 3.5 can also be obtained from Lemma 8 in [Yoshitomo Baba, Mamabu Harada, On almost M -projectives and almost M -injectives, Tsukuba J. Math. 14 (1) (1990) 53–69].

(2) Theorem 4.3 has now been extended “to solvable groups and linear groups.”

Acknowledgments

Portion of this work was done while J.B.Srivastava was visiting Ohio University–Zanesville. He would like to thank Dean James W. Fonseca for providing this opportunity. K.I. Beidar would like to express his thanks to Department of Mathematics and Center of Ring Theory and its Applications, Ohio University for providing him the opportunity to work on this project during his visit in Fall and Winter 2001–2002.

References

- [1] S. Barthwal, S.K. Jain, P. Kanwar, S.R. López-Permouth, Nonsingular semiperfect CS-rings, *J. Algebra* 203 (1998) 361–373.
- [2] A. Behn, Polycyclic group rings whose principal ideals are protective, *J. Algebra* 232 (2000) 697–707.
- [3] G.F. Birkenmeier, J.Y. Kim, J.K. Park, A counterexample for CS-rings, *Glasg. Math. J.* 42 (2) (2000) 263–269.
- [4] A.W. Chatters, C.R. Hajarnavis, Rings in which every complement right ideal is a direct summand, *Quart. J. Math.* 28 (1977) 61–80.

- [5] N.V. Dung, D. Van Huynh, P.F. Smith, R. Wisbauer, Extending Modules, Pitman, London, 1994.
- [6] J.L. Gómez Pardo, P.A. Guil Asensio, Every Σ -CS module has an indecomposable decomposition, Proc. Amer. Math. Soc. 129 (2001) 947–954.
- [7] D. Van Huynh, S.K. Jain, S.R. López-Permouth, On the symmetry of Goldie and CS conditions over prime rings, Proc. Amer. Math. Soc. 128 (2000) 3153–3157.
- [8] S.K. Jain, P. Kanwar, S.R. López-Permouth, Nonsingular semiperfect CS-rings II, Bull. London Math. Soc. 32 (2000) 421–431.
- [9] S.K. Jain, P. Kanwar, S. Malik, J.B. Srivastava, KD_∞ is a CS-algebra, Proc. Amer. Math. Soc. 128 (2) (2000) 397–400.
- [10] T.Y. Lam, Lectures on Modules and Rings, Springer, 1999.
- [11] T.Y. Lam, A First Course in Noncommutative Rings, 2nd Edition, Springer, 2001.
- [12] M.M. Parmenter, P.N. Stewart, Excellent extensions, Comm. Algebra 16 (1988) 703–713.
- [13] D.S. Passman, The Algebraic Structure of Group Rings, John Wiley, New York, 1977.
- [14] B. Stenström, Rings of Quotients, Springer-Verlag, 1975.
- [15] R. Wisbauer, Foundations of Module and Ring Theory, Gordon and Breach, Reading, UK, 1991.